

December 2000
Tashkent

Approximate formula for the ground state energy of anyons in $2D$ parabolic well

B. Abdullaev¹, M. Musakhanov¹, A. Nakamura²

¹ *Theoretical Physics Dept, Uzbekistan National University,
Tashkent 700174, Uzbekistan*

e-mail: bah_abd@iaph.silk.org, yousuf@iaph.silk.org

² *RIISE, Hiroshima University, Japan*

Abstract

We determine approximate formula for the ground state energy of anyons in $2D$ parabolic well which is valid for the arbitrary anyonic factor ν and number of particles N in the system. We assume that centre of mass motion energy is not excluded from the energy of the system. Formula for ground state energy calculated by variational principle contains logarithmic divergence at small distances between two anyons which is regularized by cut-off parameter. By equating this variational formula to the analogous formula of Wu near bosonic limit ($\nu \sim 0$) we determine the value of the cut-off and thus derive the approximate formula for the ground state energy for the any ν and N . We checked this formula at $\nu = 1$, when anyons become fermions, for the systems containing two to thirty particles. We find that our approximate formula has an accuracy within 6%. It turns out, at the big number N limit the ground state energy has square root dependence on factor ν .

I. INTRODUCTION

Anyons - two-dimensional particles obeying fractional statistics are one of possible quasi-particle excitations in fractional quantum Hall effect and High T_c superconductors [1–5].

The determination of the energy spectrum of anyons in $2D$ parabolic well is actual and interesting task of quantum mechanics. The analytical calculation of the energy spectrum of the two anyons in $2D$ parabolic well was performed in [6,7]. The energy spectrum of systems containing more than two anyons in the same external field can not be obtained analytically.

Wu [8] generalized the class of exact two anyon solution into three anyons containing system and found it's energy spectrum. There are the numerical calculations of the lowest part of the energy spectrum of three [9,10] and four [11] anyons in $2D$ parabolic well.

There is separate quantization of centre mass motion energy and relative motion energy to each other of anyons in $2D$ parabolic well [7,8]. The ground state of relative motion energy of the systems of two, three and four anyons can be linearly interpolated between bosonic and fermionic spectra [6–11] when anyonic factor ν changes from 0 to 1.

Linear dependence of full ground state energy (without separation of centre mass motion energy) from anyonic factor ν exists for arbitrary number of anyons only near the bosonic limit of spectra [8], i.e. at $\nu \rightarrow 0$. Beginning with the case of three anyons it deviates from linear character near the fermionic limit of spectra [8], i.e. at $\nu \rightarrow 1$.

By variational calculation we determine approximate formula for the ground state energy of anyons in $2D$ parabolic well which is valid for the arbitrary anyonic factor ν and number of particles N in system. We assume that centre of mass motion energy is not excluded from energy of system and anyons represented by bosonic variational wave function. As in Laughlin's variational calculation of infinite system of free anyons [12], our formula for the ground state energy calculated by variational principle contains logarithmic divergence at small distances between two anyons regularized by cut-off. We equate our variational ground state energy formula to the same formula of Wu [8] for anyons in $2D$ parabolic well near the bosonic limit on the factor ν to determine the cut-off and thus derive the approximate formula for the ground state energy at the any factor ν and number N . We check this formula at $\nu = 1$, when anyons become fermions, for the systems containing two to thirty particles. We find that our approximate formula has an accuracy within 6%. It turns out, at the big number N limit the ground state energy has square root dependence on factor ν .

The following section 2 describes the hamiltonian of the system of anyons in $2D$ parabolic well and the choice of trial wave function for the variational calculation and physical argument for this choice. Section 3 contains results of variational calculation.

II. HAMILTONIAN AND WAVE FUNCTION OF SYSTEM

The Hamiltonian of N anyons system in $2D$ parabolic well has a form:

$$\hat{H} = \frac{1}{2M} \sum_{i=1}^N (\vec{p}_i + \vec{A}(\vec{r}_i))^2 + \frac{M\omega_o^2}{2} \sum_{i=1}^N \vec{r}_i^2. \quad (1)$$

Here M - mass of particle, $\vec{p} = -i\hbar\vec{\nabla}$, where $\vec{\nabla} = \vec{i}\frac{\partial}{\partial x} + \vec{j}\frac{\partial}{\partial y}$, ω_o - eigen frequency of oscillatory motion of free particles in $2D$ parabolic well and $\vec{r}_i$ - radius vector of i - th particle number.

Vector potential for anyons $\vec{A}(\vec{r}_i)$ [8,12] in (1) is

$$\vec{A}(\vec{r}_i) = \hbar\nu \sum_{j>i}^N \frac{\vec{z} \times \vec{r}_{ij}}{|\vec{r}_{ij}|^2}. \quad (2)$$

Here $\vec{z}$ is unit vector perpendicular to $2D$ plane where situated system, $\vec{r}_{ij} = \vec{r}_i - \vec{r}_j$ and anyonic factor ν characterizes the form of fractional statistics (spin of the anyon): it can be changed from $\nu = 0$ - noninteracting bosons to $\nu = 1$ - free fermions.

In the variational consideration the ground state energy of system determined as minimum of energy

$$E = \frac{\int \Psi_T^*(\vec{R}) \hat{H} \Psi_T(\vec{R}) d\vec{R}}{\int \Psi_T^*(\vec{R}) \Psi_T(\vec{R}) d\vec{R}}. \quad (3)$$

Here $\hat{H}$ - the Hamiltonian, $\Psi_T(\vec{R})$ - trial wave function of system that has a variational parameter and $\vec{R}$ - the coordinates of all particles. Minimum energy of E (3) is reached by variation of variational parameter of trial wave function.

We suppose that trial wave function is provided by mean field approximation of Fetter, Hanna and Laughlin [13]. In this approximation particles move in the mean field $\vec{A}$:

$$\vec{A}(\vec{r}) = \frac{1}{2} \vec{B} \times \vec{r} \quad (4)$$

induced by the average density ρ of particles in the system where $\vec{B} = 2\pi\rho\hbar\nu\vec{z}$, $\rho = \frac{1}{\pi r_o^2}$ and r_o - mean distance between particles. According magnetic field $\vec{B}$ one can introduce corresponding magnetic length $a_H = (\hbar/B)^{1/2}$ and cyclotron frequency $\omega_c = B/M$. Energetic spectrum of Hamiltonian (1) with $\vec{A}$ and $2D$ parabolic well has the same energetic spectrum structure as a free particle in $2D$ parabolic well [14].

Thus assuming a bosonic representation of anyons

$$\Psi_T(\vec{R}) = \prod_{i=1}^N \Psi_T(\vec{r}_i) \quad (5)$$

for the ground state of anyons in $2D$ parabolic well one can bring wave function $\Psi_T(\vec{r}_i)$ in the following form:

$$\Psi_T(\vec{r}_i) = C \exp \left(-(\alpha + \nu) \frac{(x_i^2 + y_i^2)}{2r_o^2} \right). \quad (6)$$

Here C - normalization constant, α variational parameter and there assumed that mean distance between particles r_o equals of $R_o = (\hbar/M\omega_o)^{1/2}$ - radius localization for the wave function in the ground state of $2D$ oscillator. This equality follows from fact that interaction between two anyons at any not zero anyonic factor ν has repulsive character and relation $S_n = n\pi R_o^2$ where S_n - area that occupies by free particle at n - th energetic level of $2D$ oscillator. Last relation for the area S_n one can easily get by using a virial theorem for $2D$ oscillator.

It is convenient to express all energetic quantities by energetic quanta of $2D$ oscillator $\hbar\omega_o$ and length quantities by r_o .

Normalized trial wave function of system has a form:

$$\Psi_T(\vec{R}) = \left(\frac{\alpha + \nu}{\pi} \right)^{N/2} \prod_{i=1}^N \exp \left(-(\alpha + \nu) \frac{(x_i^2 + y_i^2)}{2} \right). \quad (7)$$

III. VARIATIONAL CONSIDERATION. RESULT OF CALCULATION.

At the calculation of variational energy E (3) it is convenient to deal with quantity $E_L(\vec{R}) = \Psi_T^{-1}(\vec{R})\hat{H}\Psi_T(\vec{R})$.

As trial wave function $\Psi_T(\vec{R})$ is not true of the Hamiltonian (1), so

$$E_L(\vec{R}) = ReE_L(\vec{R}) + iImE_L(\vec{R}). \quad (8)$$

By the acting of the Hamiltonian $\hat{H}$ (1) on $\Psi_T(\vec{R})$ (7) we find

$$ReE_L(\vec{R}) = \sum_{i=1}^N [\alpha + \nu + \frac{x_i^2 + y_i^2}{2} - \frac{(\alpha + \nu)^2}{2}(x_i^2 + y_i^2) + \frac{\nu^2}{2}(\vec{A}(\vec{r}_i))^2], \quad (9)$$

and

$$ImE_L(\vec{R}) = -\nu(\alpha + \nu) \sum_{i=1}^N (\vec{A}(\vec{r}_i)\vec{r}_i). \quad (10)$$

In the expression (10) we have a scalar product of vector $\vec{A}(\vec{r}_i)$ and $\vec{r}_i$.

A variational energy E (3) expressed by $E_L(\vec{R})$ is

$$E = \int \Psi_T(\vec{R}) E_L(\vec{R}) \Psi_T(\vec{R}) d\vec{R}. \quad (11)$$

Direct calculation gives

$$\int \Psi_T(\vec{R}) ImE_L(\vec{R}) \Psi_T(\vec{R}) d\vec{R} = 0. \quad (12)$$

So, on energy E contributes only $ReE_L(\vec{R})$.

Integrals (11) with $\Psi_T(\vec{R})$ (7) of first three terms in square brackets (9) are calculating elementary. Problems appear when we are calculating integrals with term proportional of $(\vec{A}(\vec{r}_i))^2$. This term gives two kind integrals: integrals like $\int \Psi_T(\vec{R}) \frac{1}{|\vec{r}_{ij}|^2} \Psi_T(\vec{R}) d\vec{R}$ where $i \neq j$ number of which are $N(N-1)$ and integrals like $\int \Psi_T(\vec{R}) \frac{1}{|\vec{r}_{ij}|} \frac{1}{|\vec{r}_{ik}|} \Psi_T(\vec{R}) d\vec{R}$ where $i \neq j, i \neq k$ and $j \neq k$ number of which are $N(N-1)(N-2)$.

By taking account of integrals (see Gradshtein and Ryzik [15]):

$$\int_0^\infty E_i(ax) e^{-\mu x} dx = -\frac{1}{\mu} \ln \left(\frac{\mu}{a} - 1 \right), \quad (13)$$

where constants $a > 0, Re\mu > 0$ and $\mu > a$ and

$$\int_0^\infty E_i(-\beta x) e^{-\mu x} dx = \frac{1}{\mu} \ln \left(\frac{\mu}{\beta} + 1 \right), \quad (14)$$

where constants $Re(\beta + \mu) \geq 0, \mu > 0$ and $E_i(\gamma y) = -vp \int_{-y}^\infty e^{-\gamma z} dz / z$ - the exponential integral where $\gamma > 0$ and $vp \int$ indicates integral in main quantity mean one can get

$$\int \Psi_T(\vec{R}) \frac{1}{|\vec{r}_{ij}|^2} \Psi_T(\vec{R}) d\vec{R} = (\alpha + \nu) \ln \left(\frac{2}{\delta} \right), \quad (15)$$

where cut-off parameter δ regularize the integral (15).

And an other hand the calculation gives

$$\int \Psi_T(\vec{R}) \frac{1}{|\vec{r}_{ij}|} \frac{1}{|\vec{r}_{ik}|} \Psi_T(\vec{R}) d\vec{R} = (\alpha + \nu) A, \quad (16)$$

where numerical value $A \approx 1$.

So, term by term averaged quantity $ReE_L(\vec{R})$ (9) gives expression for energy

$$E = N(\alpha + \nu) + \frac{N}{2(\alpha + \nu)} - \frac{N(\alpha + \nu)}{2} + \frac{1}{2} \nu^2 N(N-1)(\alpha + \nu) \left[\ln \left(\frac{2}{\delta} \right) - (N-2) \right]. \quad (17)$$

The extremum condition $\frac{dE}{d(\alpha + \nu)} = 0$ gives a parameter

$$(\alpha + \nu) = [1 + \nu^2(N-1) \left[\ln \left(\frac{2}{\delta} \right) - (N-2) \right]]^{-1/2} \quad (18)$$

at which minimum energy of E is

$$E_v = N[1 + \nu^2(N-1) \left[\ln \left(\frac{2}{\delta} \right) - (N-2) \right]]^{1/2}. \quad (19)$$

As established Wu [8] near the bosonic limit of ground state energy of anyons in $2D$ parabolic well on the factor ν , i.e. $\nu \rightarrow 0$, ground state energy when centre mass motion not fixed has a following expression (here and below in this section we return into normal units of energy and length):

$$E \approx \hbar\omega_o[N + N(N-1)\nu/2] \quad (20)$$

for the arbitrary number of particles N in system. So, at $\nu \rightarrow 0$ we expand expression for the energy E_v (19) on the powers ν^2 and by equating term proportional to ν^2 of expansion to second term in square brackets of (20), we find that

$$\delta = 2 \exp \left(-\frac{1 + \nu(N-2)}{\nu} \right) r_o \quad (21)$$

and finally analytical expression for the variational ground state energy is:

$$E_v = \hbar\omega_o N[1 + \nu(N-1)]^{1/2}. \quad (22)$$

At the Table 1 we bring exact energies for fermions in $2D$ parabolic well constructed by filling lowest energetic levels of $2D$ oscillator and energies calculated by formula (22). From this table one can see that deviation of variational energy E_v from exact energy E_{EXACT} is no more than 6%.

REFERENCES

- [1] F. Wilczek, Fractional Statistics and Anyon Superconductivity, World Scientific, Singapore, 1990.
- [2] S. Forte, *Rev.Mod.Phys.*, **64**, 193 (1992).
- [3] R. Iengo and K. Lechner, *Phys.Rep.*, **213**, 179 (1992).
- [4] A. Lerda, Anyons, Lecture Notes in Physics m14, Springer-Verlag, Berlin-Heidelberg, 1992.
- [5] M. Stone, editor, Quantum Hall Effect, World Scientific, Singapore, 1992.
- [6] J.M. Leinaas and J. Myrheim, *Nuovo Cimento*, **B37**, 1 (1977).
- [7] F. Wilczek, *Phys.Rev.Lett.*, **48**, 1144 (1982).
- [8] Y.-S. Wu, *Phys.Rev.Let.*, **53**, 111 (1984), Erratum *ibid* **53**, 1028 (1984).
- [9] M. Sporre, J.J.M. Verbaarschot and I. Zahed, *Phys.Rev.Lett.*, **67**, 1813 (1991).
- [10] M.V.N. Murthy, J. Law, M. Brack and R.K. Bhaduri, *Phys.Rev.Lett.*, **67**, 1817 (1991).
- [11] M. Sporre, J.J.M. Verbaarschot and I. Zahed, *Phys.Rev.* **B46**, 5738 (1992).
- [12] R.B. Laughlin, *Phys.Rev.Lett.*, **60**, 2677 (1988).
- [13] A.L. Fetter, C.B. Hanna and R.B. Laughlin, *Phys.Rev.* **B39**, 9679 (1989).
- [14] V.M. Galicki, B.M. Karnakov and V.I.Kogan, "The problems on the quantum mechanics" (in Russian), Moscow, "Nauka", 1981, 648.
- [15] I.S. Gradshteyn and I.M. Ryzik, Tables of integrals, sums, expansions and products (in Russian), Moscow, Fizmatgiz, 1100, 1962.

TABLES

TABLE I. The ground state energy of N fermions in $2D$ parabolic well.

E_{EXACT} - the energies given by exact analytical formula from [14] (in $\hbar\omega_o$ units),

E_v - the energies (in $\hbar\omega_o$ units) calculated by formula (22).

N	E_{EXACT}	E_v	$(E_v - E_{EXACT})/E_{EXACT}$ in %
2	3	2.83	6
3	5	5.20	4
4	8	8.00	0
5	11	11.18	2
6	14	14.70	5
7	18	18.52	3
8	22	22.63	3
9	26	27.00	4
10	30	31.62	5
11	35	36.48	4
12	40	41.57	4
13	45	46.87	4
14	50	52.38	5
15	55	58.09	6
16	61	64.00	5
17	67	70.09	5
18	73	76.37	5
19	79	82.82	5
20	85	89.44	5
21	91	96.23	6
22	98	103.19	5
23	105	110.30	5
24	112	117.58	5
25	119	125.00	5
26	126	132.57	5
27	133	140.30	5
28	140	148.16	6
29	148	156.17	6
30	156	164.32	5